

Self-help series fact sheet 17

Immunisation

Regular COVID-19 vaccination is recommended even in individuals who have had a past infection, and there are no safety concerns for individuals receiving a COVID-19 vaccine following a recent COVID-19 infection.

Regular COVID-19 vaccinations or boosters are the best way to protect yourself against severe illness, hospitalisation and death from COVID-19.

In particular, they are important for anyone aged 65 years or older and people at higher risk of severe COVID-19.

As with all vaccinations, people are encouraged to discuss the vaccine options available to them with their health practitioner.

The most frequently reported side effects include injection-site reactions, such as a sore arm. Other reported side effects include:

- headache
- muscle pain
- fever and chills
- fatigue.

These side effects typically last no more than a couple of days, and you will recover without any problems. If you are concerned about side effects please talk to your GP.

We encourage you to rely on credible information to make informed choices about the vaccine you receive.



This fact sheet is supported by funding from the ACT PHN through the Australian Government's PHN Program.

Funded by



Like your other vaccinations, we encourage you to discuss COVID-19 vaccine options available to you with your healthcare provider. They can help you decide how frequently you have your regular dose with an individual risk–benefit assessment.

If you need further information or support please have a look at our website at www.EmergeACT.org.au or the long COVID playlist on our YouTube channel at www.youtube.com/@arthritisact3717